


APPENDIX B

**NRCS Soils Series Report
Geologic Study
Ground Water Data
Wetland Analysis**

Soil Map—Madison County Area, Montana
(Stone Ridge Club Subdivision)



Septic Tank Absorption Fields—Madison County Area, Montana
(Stone Ridge Club Subdivision)










































Natural Resources
Conservation Service

Web Soil Survey 2.2
National Cooperative Soil Survey

6/18/2009
Page 1 of 6

Soil Map—Madison County Area, Montana
(Stone Ridge Club Subdivision)

MAP LEGEND

Area of Interest (AOI)	 Area of Interest (AOI)	 Very Stony Spot
Soils	 Soil Map Units	 Wet Spot
Special Point Features	 Blowout	 Other
	 Borrow Pit	Special Line Features
	 Clay Spot	 Gully
	 Closed Depression	 Short Steep Slope
	 Gravel Pit	 Other
	 Gravelly Spot	Political Features
	 Landfill	 Cities
	 Lava Flow	 PLSS Township and Range
	 Marsh or swamp	 PLSS Section
	 Mine or Quarry	Water Features
	 Miscellaneous Water	 Oceans
	 Perennial Water	 Streams and Canals
	 Rock Outcrop	Transportation
	 Saline Spot	 Rails
	 Sandy Spot	 Interstate Highways
	 Severely Eroded Spot	 US Routes
	 Sinkhole	 Major Roads
	 Slide or Slip	 Local Roads
	 Sodic Spot	
	 Spoil Area	
	 Stony Spot	

MAP INFORMATION

Map Scale: 1:6,630 if printed on A size (8.5" x 11") sheet.

The soil surveys that comprise your AOI were mapped at 1:24,000.

Please rely on the bar scale on each map sheet for accurate map measurements.

Source of Map: Natural Resources Conservation Service
Web Soil Survey URL: <http://websoilsurvey.nrcs.usda.gov>
Coordinate System: UTM Zone 12N NAD83

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Madison County Area, Montana
Survey Area Data: Version 11, Feb 26, 2008

Date(s) aerial images were photographed: 8/5/2005

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Madison County Area, Montana (MT636)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
17	Beaverell cobbly loam, cool, 0 to 6 percent slopes	8.5	4.1%
38	Cryaquolls, nearly level	6.4	3.1%
91	Nuley-Rock outcrop complex, 8 to 35 percent slopes	106.9	51.8%
147	Varney clay loam, 2 to 8 percent slopes	84.7	41.0%
Totals for Area of Interest		206.5	100.0%

Septic Tank Absorption Fields

Septic Tank Absorption Fields— Summary by Map Unit — Madison County Area, Montana						
Map unit symbol	Map unit name	Rating	Component name (percent)	Rating reasons (numeric values)	Acres in AOI	Percent of AOI
17	Beaverell cobbly loam, cool, 0 to 6 percent slopes	Very limited	Beaverell (90%)	Filtering capacity (1.00)	8.5	4.1%
			Beaverell, stony (4%)	Filtering capacity (1.00)		
			Rivra (3%)	Depth to saturated zone (1.00)		
				Filtering capacity (1.00)		
				Flooding (0.40)		
38	Cryaquolls, nearly level	Not rated	Cryaquolls (95%)		6.4	3.1%
			Nonhydric (5%)			
91	Nuley-Rock outcrop complex, 8 to 35 percent slopes	Very limited	Nuley (65%)	Seepage, bottom layer (1.00)	106.9	51.8%
				Slope (1.00)		
				Depth to bedrock (0.99)		
				Slow water movement (0.50)		
			Rentsac (10%)	Depth to bedrock (1.00)		
				Slope (1.00)		
				Seepage, bottom layer (1.00)		
				Large stones content (0.01)		
			Nuley, thick surface (5%)	Seepage, bottom layer (1.00)		
				Depth to bedrock (0.99)		
				Slope (0.63)		
				Slow water movement (0.50)		
147	Varney clay loam, 2 to 8 percent slopes	Somewhat limited	Varney (90%)	Slow water movement (0.50)	84.7	41.0%
			Varney, cobbly loam (7%)	Slow water movement (0.50)		
Totals for Area of Interest					206.5	100.0%

Septic Tank Absorption Fields— Summary by Rating Value		
Rating	Acres in AOI	Percent of AOI
Very limited	115.4	55.9%
Somewhat limited	84.7	41.0%
Null or Not Rated	6.4	3.1%
Totals for Area of Interest	206.5	100.0%

Engineering Properties

Madison County Area, Montana

Absence of an entry indicates that the data were not estimated. The asterisk "*" denotes the representative texture; other possible textures follow the dash.

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percent passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10 Inches	3-10 Inches	4	10	40	200		
	<i>In</i>				<i>Pct</i>	<i>Pct</i>					<i>Pct</i>	
17: Beaverell	0-4	Cobbly loam	CL-ML, ML, SC-SM, SM	A-4	0	15-30	80-100	70-90	55-80	45-70	20-30	NP-10
	4-17	Very gravelly clay loam, Very gravelly loam, Very gravelly sandy clay loam	GC, GC-GM	A-2	0	0-25	40-60	30-50	20-45	15-35	25-40	5-15
	17-60	Stratified very gravelly loamy sand	GM, GP-GM, SM, SP-SM	A-1	0	10-45	35-65	25-55	10-45	5-15	---	NP
38: Cryaquolls	---	---	---	---	---	---	---	---	---	---	---	---
91: Nuley	0-4	Sandy loam	SC-SM, SM	A-2, A-4	0	0	80-100	75-95	45-65	25-50	20-30	NP-10
	4-11	Clay loam, Sandy clay loam	CL, CL-ML, SC, SC-SM	A-4, A-6	0	0	80-100	75-95	50-75	40-65	25-40	5-15
	11-24	Gravelly sandy loam, Loam, Sandy loam	ML, SM	A-2, A-4	0	0	75-100	70-95	40-70	30-60	15-25	NP-5
	24-42	Gravelly coarse sand, Gravelly loamy coarse sand	SM, SW-SM	A-1	0	0-5	60-75	50-65	5-40	5-25	---	NP
	42-46	Unweathered bedrock	---	---	---	---	---	---	---	---	---	---
Rock outcrop	---	---	---	---	---	---	---	---	---	---	---	---

Engineering Properties

Madison County Area, Montana

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percent passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10 Inches	3-10 Inches	4	10	40	200		
	<i>In</i>				<i>Pct</i>	<i>Pct</i>					<i>Pct</i>	
91: Rentsac	0-4	Channery loam	GM, ML, SM	A-4	0	0-15	70-85	60-75	45-65	35-55	20-25	NP-5
	4-16	Very channery loam, Very flaggy loam, Extremely gravelly sandy loam	GM, SM	A-1, A-2, A-4	0	15-50	40-75	30-65	15-45	10-40	20-25	NP-5
	16-20	Unweathered bedrock	---	---	---	---	---	---	---	---	---	---
147: Varney	0-5	Clay loam	CL, CL-ML	A-4, A-6	0	0-5	90-100	80-100	60-95	55-85	25-35	5-15
	5-16	Clay loam, Gravelly clay loam, Gravelly sandy clay loam	CL, GC, SC	A-2, A-6	0	0-10	65-100	55-95	45-85	30-70	30-40	10-20
	16-48	Gravelly sandy clay loam, Gravelly sandy loam, Loam	CL, CL-ML, GC-GM, SC-SM	A-2, A-4, A-6	0	0-10	65-100	55-95	40-80	30-65	20-35	5-15
	48-60	Stratified gravelly loamy sand to loam	SM	A-1, A-2, A-4	0	0-10	70-100	55-95	35-75	20-50	15-25	NP-5

Chemical Soil Properties

Madison County Area, Montana

Map symbol and soil name	Depth	Cation- exchange capacity	Effective cation- exchange capacity	Soil reaction	Calcium carbon- ate	Gypsum	Salinity	Sodium adsorption ratio
	<i>In</i>	<i>meq/100 g</i>	<i>meq/100 g</i>	<i>pH</i>	<i>Pct</i>	<i>Pct</i>	<i>mmhos/cm</i>	
17:								
Beaverell	0-4	10-20	---	6.6-7.8	0	0	0.0	0
	4-17	15-25	---	6.6-7.8	0	0	0.0	0
	17-60	0.0-5.0	---	7.4-8.4	0-15	0	0.0-2.0	0
38:								
Cryaquolls	---	---	---	---	---	---	---	---
91:								
Nuley	0-4	10-15	---	6.6-7.8	0	0	0.0-2.0	0
	4-11	20-25	---	6.6-8.4	0	0	0.0-2.0	0
	11-24	5.0-10	---	7.4-8.4	15-30	0	0.0-2.0	0
	24-42	0.0-1.0	---	7.4-8.4	5-15	0	0.0-2.0	0
	42-46	---	---	---	---	---	---	---
Rock outcrop	---	---	---	---	---	---	---	---
Rentsac	0-4	5.0-10	---	6.6-8.4	0-5	0	0.0	0
	4-16	5.0-10	---	7.4-8.4	5-15	0	0.0-4.0	0
	16-20	---	---	---	---	---	---	---
147:								
Varney	0-5	15-20	---	6.6-7.3	0	0	0.0-2.0	0
	5-16	20-25	---	6.6-7.8	0	0	0.0-2.0	0
	16-48	10-15	---	7.4-8.4	15-30	0	0.0-2.0	0
	48-60	5.0-10	---	7.9-8.4	1-15	0	0.0-2.0	0

Physical Soil Properties

Madison County Area, Montana

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										Kw	Kf	T		
17:	<i>In</i>	<i>Pct</i>	<i>Pct</i>	<i>Pct</i>	<i>g/cc</i>	<i>micro m/sec</i>	<i>In/In</i>	<i>Pct</i>	<i>Pct</i>					
Beaverell	0-4	---	---	10-27	1.15-1.35	4.00-14.00	0.12-0.16	0.0-2.9	1.0-3.0	.20	.37	2	5	56
	4-17	---	---	20-35	1.30-1.55	4.00-14.00	0.06-0.10	3.0-5.9	0.5-1.0	.10	.32			
	17-60	---	---	0-5	1.45-1.65	141.00	0.02-0.04	0.0-2.9	0.0-0.5	.05	.15			
38:														
Cryaquolls	---	---	---	---	---	---	---	---	---	---	---	---	---	---
91:														
Nuley	0-4	---	---	15-20	1.20-1.40	4.00-14.00	0.12-0.16	0.0-2.9	2.0-4.0	.24	.24	3	3	86
	4-11	---	---	20-35	1.40-1.60	4.00-14.00	0.12-0.15	3.0-5.9	1.0-2.0	.32	.32			
	11-24	---	---	5-15	1.45-1.65	4.00-14.00	0.10-0.13	0.0-2.9	0.5-1.0	.24	.24			
	24-42	---	---	0-5	1.55-1.75	42.00-141.00	0.02-0.03	0.0-2.9	0.0-0.5	.10	.15			
	42-46	---	---	---	---	---	---	---	---	---	---			
Rock outcrop	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Rentsac	0-4	---	---	7-18	1.25-1.45	14.00-42.00	0.12-0.16	0.0-2.9	0.5-2.0	.24	.37	1	5	56
	4-16	---	---	7-18	1.30-1.55	14.00-42.00	0.07-0.09	0.0-2.9	0.5-1.0	.10	.37			
	16-20	---	---	---	---	---	---	---	---	---	---			
147:														
Varney	0-5	---	---	27-30	1.15-1.35	4.00-14.00	0.14-0.18	3.0-5.9	2.0-4.0	.28	.28	5	6	48
	5-16	---	---	30-35	1.45-1.65	4.00-14.00	0.12-0.15	3.0-5.9	1.0-2.0	.20	.32			
	16-48	---	---	10-30	1.50-1.70	4.00-14.00	0.11-0.13	0.0-2.9	0.5-1.0	.20	.20			
	48-60	---	---	5-15	1.50-1.75	4.00-14.00	0.09-0.10	0.0-2.9	0.0-0.5	.20	.20			

Source of Sand and Gravel

Madison County Area, Montana

[Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.00 to 0.99. The larger the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel]

Map symbol and soil name	Pct. of map unit	Potential as a source of gravel		Potential as a source of sand	
		Rating class and limiting features	Value	Rating class and limiting features	Value
17:					
Beaverell	90	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
38:					
Cryaquolls	95	Not rated		Not rated	
91:					
Nuley	65	Poor		Fair	
		Bottom layer	0.00	Thickest layer	0.23
		Thickest layer	0.00	Bottom layer	0.54
Rock outcrop	20	Not rated		Not rated	
Rentsac	10	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
147:					
Varney	90	Poor		Fair	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.10

Source of Reclamation Material, Roadfill, and Topsoil

Madison County Area, Montana

[Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation]

Map symbol and soil name	Pct. of map unit	Potential as a source of reclamation material		Potential as a source of roadfill		Potential as a source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
17:							
Beaverell	90	Poor		Fair		Poor	
		Droughty	0.00	Cobble content	0.72	Rock fragments	0.00
		Organic matter content low	0.13			Hard to reclaim (rock fragments)	0.00
38:							
Cryaquolls	95	Not rated		Not rated		Not rated	
91:							
Nuley	65	Poor		Fair		Poor	
		Too sandy	0.00	Depth to bedrock	0.04	Too sandy	0.00
		Droughty	0.11	Slope	0.18	Rock fragments	0.00
		Organic matter content low	0.13			Slope	0.00
		Carbonate content	0.80			Hard to reclaim (rock fragments)	0.18
Rock outcrop	20	Not rated		Not rated		Not rated	
Rentsac	10	Poor		Poor		Poor	
		Droughty	0.00	Depth to bedrock	0.00	Rock fragments	0.00
		Depth to bedrock	0.00	Cobble content	0.78	Depth to bedrock	0.00
		Organic matter content low	0.88	Slope	0.92	Slope	0.00
		Cobble content	0.99				
147:							
Varney	90	Fair		Good		Fair	
		Carbonate content	0.80			Rock fragments	0.13
		Organic matter content low	0.88				

Roads and Streets, Shallow Excavations, and Lawns and Landscaping

Madison County Area, Montana

[Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. The table shows only the top five limitations for any given soil. The soil may have additional limitations]

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
17: Beaverell	90	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Droughty Large stones content	0.96 0.84
38: Cryaquolls	95	Not rated		Not rated		Not rated	
91: Nuley	65	Very limited Slope Frost action	1.00 0.50	Very limited Cutbanks cave Slope Depth to hard bedrock	1.00 1.00 0.96	Very limited Slope Droughty	1.00 0.08
Rock outcrop	20	Not rated		Not rated		Not rated	
Rentsac	10	Very limited Depth to hard bedrock Slope Frost action Large stones content	1.00 1.00 0.50 0.01	Very limited Depth to hard bedrock Slope Cutbanks cave Large stones content	1.00 1.00 0.10 0.01	Very limited Droughty Depth to bedrock Slope Gravel content Large stones content	1.00 1.00 1.00 0.08 0.03
147: Varney	90	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	

Paths, Trails, and Golf Fairways

Madison County Area, Montana

[Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. The table shows only the top five limitations for any given soil. The soil may have additional limitations]

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
17:							
Beaverell	90	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Droughty Large stones content	0.96 0.84
38:							
Cryaquolls	95	Not rated		Not rated		Not rated	
91:							
Nuley	65	Somewhat limited Slope	0.82	Not limited		Very limited Slope Droughty	1.00 0.08
Rock outcrop	20	Not rated		Not rated		Not rated	
Rentsac	10	Somewhat limited Slope	0.08	Not limited		Very limited Droughty Depth to bedrock Slope Gravel content Large stones content	1.00 1.00 1.00 0.08 0.03
147:							
Varney	90	Not limited		Not limited		Not limited	

Sewage Disposal

Madison County Area, Montana

[Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. The table shows only the top five limitations for any given soil. The soil may have additional limitations]

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
17:					
Beaverell	90	Very limited		Very limited	
		Filtering capacity	1.00	Seepage	1.00
				Slope	0.08
38:					
Cryaquolls	95	Not rated		Not rated	
91:					
Nuley	65	Very limited		Very limited	
		Seepage, bottom layer	1.00	Slope	1.00
		Slope	1.00	Seepage	1.00
		Depth to bedrock	0.99	Depth to hard bedrock	0.96
		Slow water movement	0.50		
Rock outcrop	20	Not rated		Not rated	
Rentsac	10	Very limited		Very limited	
		Depth to bedrock	1.00	Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage, bottom layer	1.00	Seepage	1.00
		Large stones content	0.01	Large stones content	0.04
147:					
Varney	90	Somewhat limited		Somewhat limited	
		Slow water movement	0.50	Slope	0.68
				Seepage	0.50

Dwellings With Basements

Dwellings With Basements— Summary by Map Unit — Madison County Area, Montana						
Map unit symbol	Map unit name	Rating	Component name (percent)	Rating reasons (numeric values)	Acres in AOI	Percent of AOI
17	Beaverell cobbly loam, cool, 0 to 6 percent slopes	Not limited	Beaverell (90%)		8.5	4.1%
			Beaverell, stony (4%)			
			Trimad (3%)			
38	Cryaquolls, nearly level	Not rated	Cryaquolls (95%)		6.4	3.1%
			Nonhydric (5%)			
91	Nuley-Rock outcrop complex, 8 to 35 percent slopes	Very limited	Nuley (65%)	Slope (1.00)	106.9	51.8%
				Depth to hard bedrock (0.96)		
			Rentsac (10%)	Depth to hard bedrock (1.00)		
				Slope (1.00)		
				Large stones content (0.01)		
147	Varney clay loam, 2 to 8 percent slopes	Not limited	Varney (90%)		84.7	41.0%
			Varney, cobbly loam (7%)			
Totals for Area of Interest					206.5	100.0%


Dwellings With Basements— Summary by Rating Value		
Rating	Acres in AOI	Percent of AOI
Very limited	106.9	51.8%
Not limited	93.2	45.2%
Null or Not Rated	6.4	3.1%
Totals for Area of Interest	206.5	100.0%

Geologic Study



STONE RIDGE CLUB

Geologic MAP

LOCATED IN THE SE 1/4 OF SEC. 26 AND NE 1/4 OF SEC. 35,
T. 4 S., R. 2 W. OF P.M.M., MADISON COUNTY, MONTANA



Scale 1"=1/2 Mile



T4S R2W

Map Units

Qal	Alluvium of modern channels and flood plains
Qc	Colluvium
Qgr	Gravel deposit
Aqg	Quartzofeldspathic gneiss
Agph	Garnetiferous gneiss of Tobacco Root Mtns.
Ahga	Hornblende-plagioclase gneiss and amphibolite
Trl	Rhyolite flows




Engineering and Surveying Inc.

1091 Stoneridge Drive • Bozeman, MT 59718

Phone (406) 587-1115 • Fax (406) 587-9768


www.chengineers.com • info@chengineers.com

Adapted from: Montana Bureau of Mines and Technology, Geologic Map of the Ennis 30' x 60' Quadrangle, Madison & Gallatin Counties, Montana and Park County, Wyoming, 2006.




MAP UNITS	
100	Recent alluvium and flood plain
101	Recent alluvium
102	Alluvium
103	Recent alluvium
104	Recent alluvium
105	Recent alluvium
106	Recent alluvium
107	Recent alluvium
108	Recent alluvium
109	Recent alluvium
110	Recent alluvium
111	Recent alluvium
112	Recent alluvium
113	Recent alluvium
114	Recent alluvium
115	Recent alluvium
116	Recent alluvium
117	Recent alluvium
118	Recent alluvium
119	Recent alluvium
120	Recent alluvium
121	Recent alluvium
122	Recent alluvium
123	Recent alluvium
124	Recent alluvium
125	Recent alluvium
126	Recent alluvium
127	Recent alluvium
128	Recent alluvium
129	Recent alluvium
130	Recent alluvium
131	Recent alluvium
132	Recent alluvium
133	Recent alluvium
134	Recent alluvium
135	Recent alluvium
136	Recent alluvium
137	Recent alluvium
138	Recent alluvium
139	Recent alluvium
140	Recent alluvium
141	Recent alluvium
142	Recent alluvium
143	Recent alluvium
144	Recent alluvium
145	Recent alluvium
146	Recent alluvium
147	Recent alluvium
148	Recent alluvium
149	Recent alluvium
150	Recent alluvium
151	Recent alluvium
152	Recent alluvium
153	Recent alluvium
154	Recent alluvium
155	Recent alluvium
156	Recent alluvium
157	Recent alluvium
158	Recent alluvium
159	Recent alluvium
160	Recent alluvium
161	Recent alluvium
162	Recent alluvium
163	Recent alluvium
164	Recent alluvium
165	Recent alluvium
166	Recent alluvium
167	Recent alluvium
168	Recent alluvium
169	Recent alluvium
170	Recent alluvium
171	Recent alluvium
172	Recent alluvium
173	Recent alluvium
174	Recent alluvium
175	Recent alluvium
176	Recent alluvium
177	Recent alluvium
178	Recent alluvium
179	Recent alluvium
180	Recent alluvium
181	Recent alluvium
182	Recent alluvium
183	Recent alluvium
184	Recent alluvium
185	Recent alluvium
186	Recent alluvium
187	Recent alluvium
188	Recent alluvium
189	Recent alluvium
190	Recent alluvium
191	Recent alluvium
192	Recent alluvium
193	Recent alluvium
194	Recent alluvium
195	Recent alluvium
196	Recent alluvium
197	Recent alluvium
198	Recent alluvium
199	Recent alluvium
200	Recent alluvium

Base from U.S. Geological Survey
7.5-minute topographic quadrangle
Map scale 1:50,000
Projection UTM zone 12, 1983 datum



Geologic map of the Ennis 30x60 Quadrangle, Madison and Gallatin Counties, Montana and Park County, Wyoming
Karl S. Kellogg and Van S. Williams,
U.S. Geological Survey, Denver, CO



Montana Bureau of Mines and Geology
Open File 529
Geologic Map of the Ennis
30' x 60' Quadrangle, Madison and
Gallatin Counties, Montana and
Park County, Wyoming
Karl S. Kellogg and Van S. Williams,
U.S. Geological Survey, Denver, CO


GEOLOGIC MAP OF THE ENNIS 30' x 60' QUADRANGLE
MADISON AND GALLATIN COUNTIES, MONTANA, AND PARK COUNTY, WYOMING

by

Karl S. Kellogg* and Van S. Williams*

Montana Bureau of Mines and Geology
Open-File Report 529

2005



* U. S. Geological Survey, Denver, CO

Revisions by MBMG

This report has been reviewed for conformity with Montana Bureau of Mines & Geology's technical and editorial standards.

Introductory Note

This geologic quadrangle map is based almost entirely on the original map by Karl S. Kellogg and Van S. Williams (2000) published by the U.S. Geological Survey as Geologic Investigation Series I-2690, scale 1:100,000. Differences from the original map arise from MBMG's effort to integrate the Ennis quadrangle data with adjacent MBMG quadrangle maps to establish a seamless geologic coverage in southwest Montana. To this end, some map unit letter symbols have been revised, twelve new units have been added, two units have been dropped, and a few unit text descriptions have been slightly modified.





Figure 1. Location of the Ennis 1:100,000 scale quadrangle.

CORRELATION OF MAP UNITS FOR THE ENNIS QUADRANGLE



DESCRIPTION OF MAP UNITS

ENNIS 30' x 60' QUADRANGLE

* Denotes a map unit that was either (1) brought into the Ennis quadrangle from a named adjacent 30' x 60' geologic quadrangle map published by MBMG, or (2) added for edge-matching clarification.

Quaternary

- Qal** **Alluvium of modern channels and flood plains (Holocene)**—Unconsolidated silt- to boulder-size, moderately sorted to well-sorted sediments in modern flood plains, including overbank deposits. Larger clasts are moderately rounded to well rounded. Maximum thickness greater than 5 m.
- Qfp*** **Flood plain deposit (Holocene) (from Hebgen Lake quadrangle)***— Sand, silt, and clay deposited in broad, open stream valleys and in confined ephemeral stream channels. Maximum thickness unknown.
- Qc** **Colluvium (Holocene and upper Pleistocene) (Qcl of Kellogg and Williams, 2000)**— Unconsolidated to slightly indurated, mostly massive, dark-brown to light-gray-brown deposits that mantle gently to moderately sloping surfaces; sediment types are intermixed by downslope movement. Colluvium contains cobbles and pebbles derived from weathering of bedrock; loess is very fine grained sand, silt, and minor clay. Commonly contains poorly to moderately developed soil profile in upper part. Includes alluvium in small channels and sheetwash on steeper hillsides. Some small areas (as in Tobacco Root Mountains) underlain by till in valleys above 2,000 m elevation. Unmapped in many areas, particularly where deposit is thin and forms discontinuous veneer. Maximum thickness probably less than 10 m.
- Qta** **Talus deposits (Holocene and upper Pleistocene?) (Qt of Kellogg and Williams, 2000)**— Angular and subangular cobbles and boulders at base of steep valley walls or cliffs. Boulders generally as large as 2 m, although in places as large as 10 m. Locally includes minor alluvial deposits and rock-glacier deposits. Maximum thickness greater than 20 m.
- Qls** **Landslide deposits (Holocene and upper Pleistocene)**—Ranges from chaotically oriented debris to almost intact slump blocks of bedrock. About 10 m to greater than 50 m thick.
- Qrg** **Rock glacier deposits (Holocene and upper Pleistocene?) (Qr of Kellogg and Williams 2000)**—Hummocky, lobate deposits of angular boulders having a frontal slope near the angle of repose; locally active. In places, grade into and include some talus deposits (unit Qt). About 20 m thick.
- Qaf*** **Alluvial fan deposits (Holocene and upper Pleistocene) (Qf of Kellogg and Williams, 2000)***—Moderately well sorted pebble- to boulder-size gravel in fan-shaped deposits at base of mountain fronts. Includes piedmont-slope deposits and debris-flow deposits.

- Qaf1*** **Alluvial fan deposits, younger (Holocene) (from Hebgen Lake quadrangle)***—Active, small to large alluvial fans along valley margins.
- Qafc** **Deposits of Cedar Creek alluvial fan (Holocene and upper Pleistocene) (Qfc of Kellogg and Williams, 2000)**— Moderately well sorted cobble and boulder gravel that forms a prominent alluvial fan at mouth of Cedar Creek in Madison Valley. Clasts rounded to sub-rounded. Surface is characterized by braided stream pattern. Locally mantled by loess less than 1 m thick. Fan formed mainly during Pinedale (about 20-14 ka) and Bull Lake (about 140-100 ka) glaciations; deposits of both ages exposed at surface (Ritter and others, 1990); minor proximal fan development during Holocene. Modern channel of Cedar Creek incised about 10 m into fan deposits near mouth of creek, indicating that proximal fan deposits are presently eroding. As much as 150 m thick. Unit shown separately from other fan deposits due to its prominent size and detailed studies (Ritter and others, 1990)
- Qgl*** **Glacial lake deposit (Pleistocene) (from Gardiner quadrangle)***— Glacial lake deposits consisting of varved silt and clay.
- Qgt** **Glacial till (upper Pleistocene) (Qti of Kellogg and Williams, 2000)**—Unsorted, unstratified, unconsolidated, subangular to subrounded boulders in an unsorted matrix as fine as silt. Most till deposited during Pinedale (about 20-14 ka) and Bull Lake (about 140-100 ka) glaciations. Pinedale-age till preserved in hummocky deposits that contain numerous closed depressions and have a thin or non-existent soil profile; deposits of Bull Lake-age till have more rounded topography, are more dissected, and generally exhibit a well-developed soil profile. As much as about 50 m thick.
- Qd** **Diamicton (upper? Pleistocene)**—Unsorted, unstratified, nonconsolidated deposit composed of angular to subangular clasts of Archean gneiss as large as 2 m across. Believed to be mostly debris-flow deposits, but may include some till and landslide deposits. Mapped only in Jack Creek area. Maximum thickness greater than 20 m.
- Qgr*** **Gravel deposits (Pleistocene) (from Hebgen Lake quadrangle; part of Qg unit of Kellogg and Williams, 2000)***--Sand and pebble- to boulder-size gravel deposit underlying several terrace levels. Located primarily along the Ruby River at elevations of as much as 100 m. In southwest corner of Ennis quadrangle map area.
- Qat*** **Terrace-gravel deposits (upper Pleistocene) (part of Qg of Kellogg and Williams, 2000)***—Moderately sorted, moderately rounded to well-rounded sand and gravel. Underlies about 12 recognized terrace surfaces in Madison Valley. Bearizi (1987) recognized 11 geomorphic surfaces along Jack Creek. In Madison Valley, east of Madison River, includes all but the highest (and oldest) terrace-gravel deposit, which is shown separately (unit Qgc). Mantled by less than 2 m of loess at most places, although loess on many higher surfaces is thick enough to support cultivation. Mostly less than 10 m thick.
- Qatc** **Terrace-gravel deposits of Cameron Bench (Pleistocene) (Qgc of Kellogg and Williams, 2000)**—Moderately sorted, moderately rounded to well-rounded sand and gravel underlying topographically highest (and oldest) terrace deposits in Madison Valley. Formed during Bull Lake glaciation (about 140-100 ka) (Schneider and Ritter, 1987) Mantled by less than 2 m of loess in most places.

Quaternary and Tertiary

QTgr* **Gravel (Pleistocene and/or Pliocene) (from Bozeman quadrangle)***—Well-rounded, moderately well-sorted, clast-supported, calcite-cemented to unconsolidated cobble and pebble gravel with some small boulders ... (Vuke and others, 2002). One small outcrop on central north border with Bozeman quadrangle.

QTbf **Basin-fill deposit (Pleistocene and Pliocene?)**—Interbedded, moderately sorted, poorly consolidated to unconsolidated, well-bedded to massive silt, sand, and well-rounded cobbles that are exposed in slopes adjacent to, and below, terraces in Madison Valley. Exposed basin-fill deposits are interpreted to be no older than Pliocene, although they overlie buried basin-fill deposits, as thick as 4,500 m, that may be as old as Eocene in the Madison Valley (Rasmussen and Fields, 1983).

Tertiary

Thr **Huckleberry Ridge Tuff (Pliocene) (Tht of Kellogg and Williams, 2000)**—Light-brown to gray, massive welded tuff that contains sparse phenocrysts of sanidine, quartz, and plagioclase. Matrix is mostly devitrified glass shards, opaque minerals, and aphanitic minerals. Contains sparse pumice fragments. Lithophysae and vesicles not observed in quadrangle. Forms prominent cap to many ridges and buttes. Age of unit is 2.0 Ma and source is from Yellowstone caldera (Christiansen and Blank, 1972), the northern edge of which is about 30 km south of quadrangle's southeastern corner. Thickness increases to south, and maximum thickness in quadrangle is about 50 m.

Tls **Landslide deposit (Pliocene and Miocene)**—Mapped at several localities in northern Madison River Valley where significant erosion and dissection has occurred since landslide events (Kellogg, 1992, 1993b).

Ts **Sediment or sedimentary rocks, undivided (Miocene)**—White, light gray, and light-brownish-gray, moderately indurated to well-indurated, locally vuggy limestone, limy sandstone, ash-bearing sandstone, pebbly sandstone, and pebble conglomerate. Interlayered ash deposits are common, especially in association with limestone. Limestone may be either lacustrine deposits or carbon ate paleosols (Hanneman and others, 1991). Mapped along the Madison River, east of Ennis Lake, and in Spanish Creek basin. One maximum $^{40}\text{Ar}/^{39}\text{Ar}$ age on sanidine from just east of Ennis Lake in Madison Valley (NW1/4 Sec. 6, T. 5 S., R. 1 E), is 16.2 ± 0.19 Ma (K, K. Kellogg and S. S. Harlan, unpub. Data, 1990). Deposits in Spanish Creek basin are undated, but probably Miocene. Maximum thickness greater than 1,000 m.

Tli **Freshwater limestone (Miocene) (Tl of Kellogg and Williams, 2000)**—Light-brownish-gray to yellowish-gray, locally vuggy, well-bedded, fine-grained limestone. Similar to thin limestone beds associated with ash-bearing sediments in unit Ts, although much thicker, so is considered to be Miocene in age; maximum thickness greater than 20 m. Mapped along west side of Madison Valley (Hadley, 1969a, b).

Volcanic rocks of Virginia City volcanic field (Oligocene and Eocene)

- Tvd** **Diatreme (Oligocene?)**—Probably basaltic, contains fragments of clay and basalt (Wier, 1982). Intrudes both volcanic rocks and Archean gneiss. One oval outcrop mapped, about 200 m in diameter, just south of Virginia City.
- Tanb** **Andesite and basalt flows (Oligocene and Eocene) (Tvv and Tv of Kellogg and Williams, 2000)**—Mostly dense to scoriaceous basalt and andesite flows, volcanic agglomerates, sandy tuff, and interbedded basaltic cinder deposits. Flows are black and dark brown and contain small phenocrysts of calcic plagioclase, olivine, augite, opaque minerals, and, in places, potassium feldspar and biotite (Hadley, 1980). At least one silicic (latitic?) Plagioclase-pyrrhic flow occurs near base of unit, 4 km southeast of Virginia City. Three localities gave the following potassium-argon (K-Ar) dates: (1) 49.3 ± 2.5 and 51.1 ± 1.9 Ma (biotite) and 51.0 ± 3.8 Ma (plagioclase); this flow is just north of Virginia City and overlies Archean basement; (2) 34.4 ± 3.0 Ma (whole rock); and (3) 32.7 ± 1.4 Ma (whole rock) (Marvin and others, 1974). Unit greater than 200 m thick. Unit includes outcrops of andesite and basalt flows (Oligocene and/or Eocene) in Madison Valley in southern part of quadrangle (Hadley, 1969b) that may be outliers of volcanic rocks of Virginia City volcanic field (see Kellogg and Williams (2000).
- Trf** **Rhyolite flows (Oligocene or Eocene) (Tvr of Kellogg and Williams, 2000)**—Dark-gray to black aphanitic rhyolite flows north and northwest of Ennis Lake (Kellogg, 1993b; Vitaliano and Cordua, 1979). Cut by many faults; local steep dips. Correlated with volcanic rocks of Virginia City volcanic field.
- Tfet** **Felsic tuff (Eocene) (Tvt of Kellogg and Williams, 2000)**—Pale-gray to white rhyolitic air-fall ash, tuffaceous sandstone, and sparse gravel lenses. Poorly exposed in most places and occurs locally at base of sequence of volcanic rocks of Virginia City volcanic field. Large areas mapped by Hadley (1969b) as felsic tuff are underlain mostly by chaotic, locally tuffaceous landslide deposits. Thickness as much as about 40 m.
- Tvs** **Volcaniclastic sandstone and gravel deposits (Eocene) (Tg of Kellogg and Williams, 2000)**—Coarse-grained, well-bedded, cross-bedded volcaniclastic sandstone, pebbly sandstone, and pebble and cobble conglomerate that in places underlie the rocks of the Virginia City volcanic field (Hadley, 1969b, 1980).

Volcanic rocks of undetermined affinity (Oligocene and/or Eocene)

- Tda** **Dacite (Eocene) (Ti of Kellogg and Williams, 2000)**—Light- to medium-gray and brownish-gray, dense, porphyritic dacite. Phenocrysts are euhedral plagioclase, hornblende, and biotite. Petrographically very similar to dacite porphyry of Fan and Lone mountains, but at least some stocks intrude Eocene volcanic rocks; other nearby rocks of similar composition intrude Cambrian strata and Archean gneiss. One potassium-argon biotite date of 49.5 ± 1.5 Ma (Chadwick, 1969).

Absaroka Volcanics Supergroup (Eocene)

Tia **Intrusive rocks of the Absaroka Volcanics (Eocene) (Tai of Kellogg and Williams, 2000)**—Dark-gray, very fine grained basalt or andesite intrusive breccia containing sparse, small phenocrysts of plagioclase. Forms two small necks in upper Porcupine Creek, east of Gallatin River Valley. Intrudes Eocene volcanic rocks and Archean gneiss. Not included in Absaroka Volcanic Supergroup by Simons and others (1985), but unit probably formed feeder vents for younger rocks of Supergroup.

Tav* **Absaroka Supergroup volcanic rocks, undivided (Eocene) (Tam and Tas of Kellogg and Williams, 2000)***

Mount Wallace Formation (Tam) (Description modified from Kellogg and Williams, 2000). Shown where proportion of mostly andesitic and basaltic flows predominates over volcanoclastic rocks, as mapped by the U.S. Geological Survey (1972). Compositions similar to those in Sepulcher Formation, and conformably overlies Sepulcher Formation.
Equivalent to Tanf = Andesite flows of Hyalite Peak Volcanics to east on adjacent Gardiner quadrangle.

Sepulcher Formation (Tas) (Description modified from Kellogg and Williams, 2000). Mafic and intermediate-composition volcanoclastic rocks, as mapped by the U.S. Geological Survey (1972). Most of unit composed of dark-grayish-brown, laharic breccias, volcanoclastic conglomerate, and light- to medium-gray volcanoclastic sandstone. Both laharic deposits (vent facies) and conglomeratic deposits (alluvial facies) are greater than 100 m thick in places, crudely stratified, and well indurated. Both lahars and conglomerates contain much silicified wood, commonly as large tree trunks as thick as 3 m, in growth position.
Equivalent to Tae = Andesite epiclastics of Hyalite Peak Volcanics to east on adjacent Gardiner quadrangle.

Tanf **Andesite flows of Hyalite Peak Volcanics (Eocene) (from Gardiner quadrangle)**—Commonly autobrecciated; includes some epiclastic lenses. This unit is thought to be correlative with the Mount Wallace Formation of the Sunlight Group as mapped by the U.S. Geological Survey (1972) near the northern boundary of Yellowstone National Park (After Chadwick, 1982). On Ennis quadrangle, this unit occurs as one small area at south end of east quadrangle boundary.

Tcsi **Basal conglomerate and siltstone (lower Eocene) (Tc of Kellogg and Williams, 2000)**—Cobbles and boulders predominantly of Archean rocks, with subordinate clasts of Paleozoic carbonate and Tertiary volcanic rocks, mapped near Garnet Mountain in northeastern part of quadrangle (McMannis and Chadwick, 1964). Conglomerate is locally overlain by a thin, discontinuous siltstone of late early Eocene age. Not overlain by volcanic rocks in quadrangle, but similar rocks in adjacent areas occur at the base of volcanic sequences of the Absaroka Volcanic Supergroup.

Tertiary and Cretaceous

TKda **Dacite (Eocene or Upper Cretaceous)**—(from Robinson, 1963) Medium-light-gray with local yellowish-brown or yellowish-green splotches from oxidation of iron-

bearing minerals; uniformly fine-grained. Dominantly dacite, but includes some quartz latite. Exposed thickness as much as 250 ft.

TKfi **Felsic intrusive rocks (lower Tertiary or Upper Cretaceous) (TKf of Kellogg and Williams, 2000)**—Greenish-gray to pinkish-gray, sparsely porphyritic massive, siliceous dacite or rhyolite. Most phenocrysts consist of as much as 15 percent white, strongly saussuritized plagioclase crystals as long as 3 mm. Rock commonly stained black by manganese oxide. Mainly forms thin sills, dikes, and irregular pods intruding exclusively Archean rocks. Described by Kellogg (1993b). Unit undated, but shares compositional and textural similarities with both Upper Cretaceous dacite porphyry of Fan and Lone Mountains and Eocene rhyolite and rhyodacite plugs near Norris (Kellogg, 1994, 1995), about 10-15 km north of quadrangle.

TKga **Gabbro sills (Eocene to Upper Cretaceous) (TKg of Kellogg and Williams, 2000)**—Black, medium-grained plagioclase-clinopyroxene gabbro that typically weathers into spheroidal blocks. In places completely weathered into grussy orange-brown soil. Several larger sills contain irregular dikes, as wide as 2 m, of fine-grained, pinkish-gray, inequigranular clinopyroxene-biotite syenite, in which the clinopyroxene forms conspicuous rods as long as 1 cm. Syenite, in turn, is intruded by thin aplitic dikes. Unit crops out near Meadow Village at Big Sky and intrudes Upper and Lower Cretaceous rocks.

Cretaceous

Kdap **Dacite porphyry of Fan and Lone Mountains (Upper Cretaceous) (Kd of Kellogg and Williams, 2000)**—Gray to greenish-gray porphyritic dacite that weathers to a very light gray or tan. Euhedral to subhedral phenocrysts compose 30-50 percent of the rock; phenocrysts are 70-90 percent zoned plagioclase crystals (An_{25-30}) as long as 2 cm, trace to 15 percent green hornblende crystals as long as 6 mm, 0-15 percent biotite flakes as long as 3 mm, and 0-10 percent equant quartz crystals as long as 5 mm. Mafic minerals are commonly altered to chlorite and epidote; plagioclase is sericitized. Matrix is very fine grained and dense, and contains about 5 percent opaque minerals. Fine-grained chill zone extends inward several centimeters from contacts. Commonly contains mafic autoliths as large as about 0.5 m. Forms sills, some greater than 80 m thick, in two intrusive centers underlying Fan Mountain and Lone Mountain. The sills are thought to form a “Christmas-tree laccolith” complex underlying both mountains (Swanson, 1950), although Kellogg (1992) found no evidence for a central “truck.” Late Cretaceous beds dip away from the central peaks at both Fan and Lone Mountains. Intrudes rocks that range in age from Middle Cambrian to Late Cretaceous. Where less than about 20 m thick, unit is indicated on map by a single line. Potassium-argon (K-Ar) and $^{40}Ar/^{39}Ar$ ages from hornblende are about 68-69 Ma (Tysdal and others, 1986; K.S. Kellogg and S.S. Harlan, unpublished data, 1994).

Kit **Intrusive rock of Tobacco Root Batholith, undivided (Upper Cretaceous) (Kgd of Kellogg and Williams, 2000)**—Gray, coarse-grained, inequigranular to porphyritic, massive hornblende-biotite granodiorite, monzogranite, and monzodiorite (classification of Streckeisen, 1975). Typically contains 50 percent normally zoned oligoclase ($An_{25}-An_{28}$), 15-25 percent green hornblende, 10-20 percent microcline (slight development of braid perthite) commonly as phenocrysts as long as 3 cm, 0-20 percent quartz, 5 percent biotite, 0-1 percent clinopyroxene cores in some hornblende, trace to 3 percent magnetite, and traces of apatite and

conspicuous sphene and zircon; hypidiomorphic texture. Includes minor dark-gray hornblende diorite and granodiorite along border and in satellitic stocks east of main batholith. Weathers light gray, in rounded tors or flat grussy outcrops. K-Ar age of batholith is 71-74 Ma (Vitaliano and Cordua, 1979).

- Ksc** **Sphinx Conglomerate (Upper Cretaceous) (Ks of Kellogg and Williams, 2000)**—Reddish-orange conglomerate with sandstone matrix cemented by calcite and hematite. Contains well-rounded cobbles and boulders derived from Paleozoic and Mesozoic strata shed from thrust plates to west; represents an unroofing sequence, the clasts generally increasing in age upwards (DeCelles and others, 1987). Lower contact is conformable with underlying Livingston Formation. Well exposed only on Sphinx Mountain and The Helmet. Minimum thickness about 610 m.
- Kbe** **Beaverhead Conglomerate (Upper Cretaceous) (Kb of Kellogg and Williams, 2000)**—Conglomerate and interbedded limestone containing well-rounded clasts of Archean gneiss and Belt Supergroup quartzite. Mapped by Hadley (1969b) in gravelly Range as two separate units (Sphinx Conglomerate and Tertiary gravel), but both units were reinterpreted on fossil and stratigraphic evidence as Beaverhead Group rocks of Cretaceous age (J.M. O'Neill, oral commun., 1997).
- Kbel*** **Beaverhead and Livingston Groups, undivided (Upper Cretaceous) (from Hebgen Lake quadrangle)***
- Klv** **Livingston Formation (Upper Cretaceous) (Kl of Kellogg and Williams, 2000)**—Livingston formation exposed on west side of Madison Range, near Sphinx Mountain, and east of Gallatin River, south of the Spanish Peaks fault. Does not crop out west of Madison River Valley. Undivided unit shown only in southeastern part of quadrangle. Description after Tysdal (1990).
- Klvu** **Upper member, informal, of Livingston Formation (Klu of Kellogg and Williams, 2000)**—Cobble and boulder conglomerate composed of well-rounded volcanic clasts in matrix of coarse-grained volcanoclastic sandstone; upper part is predominantly sandstone. Lower contact is conformable. Thickness as much as 180 m.
- Klvm** **Middle member, informal, of Livingston Formation (Klm of Kellogg and Williams, 2000)**—Brown, maroon, and gray dacite to basalt flows, autoclastic breccia, tuff breccia, and welded tuff; minor interlayered volcanoclastic sandstone. Lower contact is conformable. Thickness estimated 300-450 m.
- Klvi** **Lower member, informal, of Livingston Formation (Kli of Kellogg and Williams, 2000)**—Complexly intertonguing units of dark-green to dark-brown, medium- to coarse-grained, locally pebbly volcanoclastic sandstone, olivine basalt, mudflow breccia, volcanoclastic conglomerate, and mudstone. Conformable with underlying Everts(?) Formation. Thickness ranges from about 50 to 210 m).

- Kevm** **Everts Formation through Mowry Formation, undivided (Upper Cretaceous) (Ku of Kellogg and Williams, 2000)**—May locally include Muddy Sandstone and Thermopolis Shale. Mapped in area of Cretaceous outcrops east of Gallatin River.
- Kevv** **Everts Formation and Virgelle Sandstone, undivided (Upper Cretaceous) (Kev of Kellogg and Williams, 2000)**
- Everts Formation:** Thin- to medium-bedded, light-gray to dark-gray, poorly to moderately sorted, quartz-rich sandstone; intercalated with thin-bedded, greenish-gray to dark-gray mudstone and siltstone. Contains a few thin beds of dense limestone, porcellanite, and coal. About equal quantities of sandstone and finer-grained sedimentary rocks. Does not crop out west of Madison River Valley. Conformably overlies Virgelle Sandstone.
- Virgelle Sandstone:** Thin-to thick-bedded, medium- to coarse-grained, light-brown to yellowish brown, trough-cross-bedded quartz sandstone that forms prominent white-weathering ledges. Conformably overlies Telegraph Creek Formation. Does not crop out west of Madison Valley. About 25 to 50 m thick.
- Ktc** **Telegraph Creek Formation (Upper Cretaceous)**—Light-gray to dark-brown, thin-bedded to medium-bedded, feldspathic, calcite-cemented sandstone and interbedded dark-gray siltstone and mudstone. Middle marker sequence is a 20-m-thick white tuffaceous siltstone and sandstone. Contains thin flaggy sandstone beds in lower part. Conformably overlies Cody Shale. Does not crop out west of Madison River Valley. Thickness about 200 m.
- Kco** **Cody Shale (Upper Cretaceous) (Kc of Kellogg and Williams, 2000)**—Upper part consists of thin-bedded black fissile shale that is interbedded with minor amounts of thin-bedded, brown, commonly bioturbated calcareous, fine-grained sandstone. Lower part consists of black, fissile shale and minor siltstone that weathers dark gray. Does not crop out west of Madison River Valley. Conformably overlies Frontier Formation. Thickness about 300 m.
- Kf** **Frontier Formation (Upper Cretaceous)**—Mostly a sequence of alternating black shale and light-gray to yellowish-tan, thin-bedded to very thick bedded, cross-bedded sandstone. In the Madison Range, sandstone ledges are as thick as 3 m; ratio of sandstone to shale is about 1:3. Shale is locally carbonaceous or coaly. Shale sequences are as thick as about 20 m and commonly contain equally spaced 5- to 10-cm-thick sandstone beds, spaced 10-20 cm apart. Contains several white porcellanite beds; one prominent porcellanite bed in the Madison Range, about 5 m thick, is about 15 m above base of sequence and displays well-developed ball-and-pillow structures. Conformable with underlying Mowry Shale. Formation thickens greatly to the west; thickness in Madison Range about 140-180 m while that in Gravelly range is greater than 1,500 m.
- Km** **Mowry Formation (Upper Cretaceous) (Kmo of Kellogg and Williams, 2000)**—Mostly brownish-gray and greenish-gray tuffaceous mudstone and shale and, in upper part, contains abundant thin sandstone beds. Lower Vaughn Member consists of conspicuous gray, green, yellow, brown, orange, pink, and cream-colored bentonitic mudstone, porcellanite, siltstone, and minor interbedded quartz sandstone, although strikingly varicolored units are missing in northern Madison Range. Poorly exposed in most places. Unconformable with underlying Muddy formation. Thickness 90-180 m.

Kmdt**Muddy Sandstone and Thermopolis Formation, undivided (Lower Cretaceous) (Kmt of Kellogg and Williams, 2000)**

Muddy Sandstone: Thin- to medium-bedded, medium- to coarse-grained, brown to brownish gray, poorly to moderately indurated, clayey, ledge-forming salt-and-pepper sandstone; locally contains mud chips as long as 1 cm. In northern Madison Range, formation typically exposed as upper and lower sandstone sequence interrupted by central, poorly exposed shaly sequence. Thickness varies widely; in the northern Madison Range it is about 20-45 m thick; farther south in the Madison Range, it is as great as 107 m (Tysdal, 1990); in the Gravelly Range it is about 15 m thick (Hadley, 1969b, 1980).

Thermopolis Formation: Composed of an upper sequence (three-quarters of the total thickness) that is black to dark-gray, locally carbonaceous, fissile shale and poorly indurated, thin-bedded, silty brown sandstone. Lower one-quarter of unit is thin- to medium-bedded, fine-grained, white to tan quartz arenite that contains black shale interbeds; ripple marks and Liesegang bands are common in sandstone beds. Unconformable contact with underlying Kootenai Formation. Thickness throughout quadrangle about 70 to 80 m.

Kk

Kootenai Formation (Lower Cretaceous)—Upper 10-15 m are medium-bedded light-gray, micritic, oolitic limestone that contains abundant gastropod fossils in uppermost part. Middle part is variegated red, purple, yellow, and gray shale, mudstone, siltstone, sandstone, and locally nodular freshwater limestone. Formation typically weathers to a reddish soil. Lower 10-40 m is medium- to coarse-grained, gray, well-indurated, ledge-forming salt-and-pepper sandstone that contains a basal chert-pebble conglomerate as thick as 1 m. Lower contact is unconformable. Total thickness in the northern Madison Range is about 90 m (Kellogg, 1992, 1993a); in the Gravelly Range it is as thick as 170 m (Hadley, 1980), and in the southeastern part of the map area it is as much as about 130 m thick.

Jurassic**Jm**

Morrison Formation (Upper Jurassic)—Upper 20-30 m is mostly black and locally purple shale that contains minor intercalated thin- to medium-bedded, rusty-brown and gray quartz sandstone that is lensoidal in places. Lower, thicker part of the Morrison is composed of thin-bedded, gray, yellow, orange, red, and green shale and siltstone interbedded with lesser amounts of gray quartz arenite and thin-bedded, brown limestone. Unconformable with underlying unit. About 75-100 m thick.

Je*

Ellis Group (Upper and Middle Jurassic)* Mapped only along south-central border with adjacent Hebgen Lake quadrangle.

Swift Formation: Brown, medium- to coarse-grained glauconitic sandstone, locally containing abundant chert pebbles, and minor olive-green shale; glauconite commonly weathered to orange limonitic clots. More calcareous in eastern part of map area (Tysdal, 1990), Unconformably overlies Rierdon Limestone. Less than 12 m thick.

Rierdon Formation: Thin- to thick-bedded, yellowish-gray to brownish-gray, fine-grained oolitic limestone; contains a few limey siltstone interbeds. Bivalves *Camptonectes* sp. and

Gryphea sp. Locally abundant. Forms prominent cliffs near Shell Creek in western Madison Range. Thickness about 8 to 30 m.

Sawtooth Formation: Interbedded limestone and shale; weathers yellowish-brown and is very poorly exposed. Unconformably overlies Dinwoody Formation. Thickness 25 to 55 m.

Jurassic and Triassic

J@ md Morrison through Dinwoody Formations, undivided (Upper and Middle Jurassic and Lower Triassic) (J@ u of Kellogg and Williams, 2000)

J@ ed Ellis Group and Woodside and Dinwoody Formations, undivided (Upper and Middle Jurassic and Lower Triassic)—In most places this combined unit is poorly exposed. Total thickness about 185 m.

Triassic

@ w* Woodside Formation (Lower Triassic; one small outcrop area extended from adjacent Hebgen Lake quadrangle)*—Brick-red to orange-red, thin-bedded siltstone and mudstone interbedded with gypsum and scattered thin beds of limestone. Mostly poorly exposed, although forms prominent red soil. Thickness increases markedly to southeast; nonexistent north of Porcupine Creek in Gallatin Range (Hadley, 1980). Unit previously mapped as Chugwater Formation in southeastern part of quadrangle (Simons and others, 1985). Thickness 0 to about 220 m (Tysdal 1990).

Dinwoody Formation (Lower Triassic; not mapped separately; combined with Woodside Formation and Ellis Group as unit J@ ed)—Brown silty sandstone, siltstone, and thin-bedded brown limestone. Small brachiopods locally abundant. Weathers light yellowish-brown and is poorly exposed. Unconformably overlies Shedhorn Sandstone. Thickness ranges from about 20 to 80 m.

Permian

Psh Shedhorn Sandstone (Lower Permian) (Ps of Kellogg and Williams, 2000)—Mostly medium- to massive-bedded, gray, fine- to coarse-grained, very well indurated quartz-rich sandstone that contains white cherty stringers and nodules. Thin, shaly phosphorite beds in middle of formation are rarely exposed. Dolomitic near base. In Gravelly Range, chert is locally massive and filled with voids, so that chert outcrops commonly have a rubbly appearance. Conformably overlies Quadrant Sandstone. Formation about 5-70 m thick, though is about 50 m thick in northern Madison Range.

Permian, Pennsylvanian, and Mississippian

PMsa Shedhorn Sandstone through Amsden Formation, undivided (Lower Permian, Pennsylvanian, and Upper Mississippian) (Pmu of Kellogg and Williams, 2000)—Mapped in Garnet Mountain quadrangle by McMannis and Chadwick (1964), who used the term "Phosphoria Formation" instead of "Shedhorn Sandstone." Units described below.

Pennsylvanian and Mississippian

IP Mqa **Quadrant and Amsden Formations, undivided (Pennsylvanian and Upper Mississippian) (&qa of Kellogg and Williams, 2000)**

Quadrant Sandstone (Pennsylvanian)— Medium- to thick-bedded , white to yellowish-tan, well-sorted, fine- to medium-grained, dolomite-cemented quartz arenite; cross beds common. Lower half and at least upper 15 m of formation contain a few medium-bedded, light-yellowish-tan dolostone beds. Crops out prominently Conformable contact with underlying Amsden Group. Thickness about 75 m.

Amsden Group (Lower Pennsylvanian and Upper Mississippian) – Varies widely, both in stratigraphy and thickness. In central and southern Madison Range, and in Gravelly Range east of Greenhorn Thrust, unit consists of about 12 to 15cm of brick-red, reddish-brown, and pink calcareous siltstone, silty shale,, shale, sandstone, and light-gray limestone and yellowish-gray dolomite. Along western Madison Range, upper part contains medium- to thin-bedded, gray to grayish-tan dolostone, locally quartzitic, that grades downward into gray limestone beds (correlated by Kellogg, 1992, with Middle and Lower Pennsylvanian Devils Pocket Formation and Alaska Bench Limestone of Wardlaw and Pecora, 1985). Total thickness of Amsden Group in western Madison Range is 88 m.

Snowcrest Range Group (Upper Mississippian)—Mapped [Observed] only in western Madison Range (Kellogg, 1992), although rocks of the group may have been either unrecognized elsewhere or placed in Amsden Group. Upper 55 m of Snowcrest Range Group (Upper Mississippian Lombard Limestone) is mostly thin- to medium-bedded, gray limestone that becomes more dolomitic and shaly toward base. The Lombard overlies Kibby Sandstone, which consists of about 75 m of thin- to medium-bedded, yellowish-gray to maroon, friable, dolomitic sandstone, sandy dolostone, and siltstone. [Note: this unit does not appear on Kellogg and Williams' (2000) map or correlation chart, though the unit is described in the text. Nor does the unit appear on this current map]

Mississippian

Mbs **Big Snowy Group, undivided (Upper Mississippian)** —Mapped only west of Greenhorn thrust in Gravelly range (Hadley, 1969b, 1980); upper, thicker part consists of thin- to thick-bedded, pale olive gray, locally very fossiliferous limestone and minor dark-greenish or olive-gray mudstone and shale. Lower 60 m or so consists of red and greenish-gray shale and calcareous sandstone and minor greenish-gray cherty limestone; sandstone and red color increase downward in sequence. Detailed description of Big Snowy Group is given by Hadley (1980). Total thickness 135-275 m.

Mm **Madison Group, undivided (Upper and Lower Mississippian)**

Mmc **Mission Canyon Limestone (Upper and Lower Mississippian)**—Medium- to massive-bedded, light-gray, gray, and brownish-gray, medium-crystalline limestone and minor dolostone. Weathers light gray. Chert stringers and nodules are common; locally fossiliferous; solution breccias in uppermost part. Prominent ridge-forming unit. Conformable with underlying Lodgepole Limestone. Thickness about 240 m.

MI

Lodgepole Limestone (Lower Mississippian)—Thin- to medium-bedded, gray to brownish-gray, finely crystalline limestone that commonly grades into silty limestone interbeds several centimeters thick. Locally cherty; upper half is profusely fossiliferous. Conformable with underlying Three Forks formation (erroneously described as unconformable above the Three Forks in Kellogg, 1992, 1993a). Thickness about 180 m.

Mississippian and Devonian

MDtj **Three Forks Formation (Lower Mississippian and Upper Devonian) and Jefferson Formation (Upper Devonian), undivided**

Three Forks Formation: Mostly thin-bedded, yellowish-orange to yellowish-tan siltstone and silty limestone containing a few thin interbeds of brick-red-weathering siltstone. Weathers light yellowish tan. Contains a medium-gray irregularly bedded, approximately 12-m-thick, rough-weathering limestone about 25 m above base of unit (Logan Gulch Member). Formation Poorly exposed; forms slopes and swales. Unconformably overlies Jefferson Formation. Thickness 40-60 m.

Jefferson Formation : Thin- to thick-bedded, black, brown, dark-gray, and light-gray, petroliferous, medium-crystalline to coarsely crystalline dolostone; colors vary considerably over short stratigraphic intervals. Weathers mostly brown, and outcrops are typically knobby and irregular; forms conspicuous brown hoodoos. At least upper 15 m is thick- to massive-bedded, gray dolostone solution breccia (Bridbear Member). Unconformably overlies Bighorn Dolomite (?) In northern Madison Range. Thickness about 110 m.

Ordovician and Cambrian

O€bp **Bighorn Dolomite through Park Shale, undivided (Ordovician through Middle Cambrian) (O€rm of Kellogg and Williams, 2000)**

Bighorn Dolomite: Medium-gray, sugary dolostone in 0.2- to 1.0-m-thick beds; weathers very light gray. Tentatively placed in Big horn Dolomite by Hanson (1952) along ridge in NE¼, sec. 22 and NW¼, sec. 23, T. 5 S., R. 1 E., in northwestern Madison Range; not recognized south of this location. Unconformably overlies Red Lion Formation. Thickness about 3 m at location noted above. Mapped east of Gallatin River and north of Spanish Peaks fault (McMannis and Chadwick, 1964).

Red Lion Formation (Upper Cambrian): Thin-bedded medium-gray to tan, siliceous dolomite containing conspicuous orange-tan to reddish-tan, cherty stringers as thick as 2cm; lower 7 m contain intraformational clasts as large as 5 mm. Along west side of Gallatin Range, contains a lower greenish-gray, locally quartzitic shale sequence as thick as 10 m (Dry Creek Shale member) (McMannis and Chadwick, 1964). Unconformably overlies Pilgrim Formation. Thickness about 40 to 60 m.

Pilgrim Dolomite (Upper Cambrian): Gray, light-gray, and brownish-gray, medium- to massive-bedded, locally oolitic, medium-crystalline dolomite. Weathers light gray and

contains irregularly shaped darker gray mottles. Conformably overlies Park Shale. Forms conspicuous crags. Thicknesses about 30 m in northern Madison Range (Kellogg, 1992), about 60 m in the south-central part of the quadrangle (Tysdal, 1990), as much as 120 m in the Gravelly Range, although missing along east side of range (Hadley, 1969b), and as much as 75 m in the northern Gallatin Range (McMannis and Chadwick, 1964).

Park Shale (Middle Cambrian): Greenish-gray to tan, fissile shale. Poorly exposed, slope-forming unit conformably overlies Meagher Limestone. Thickness about 30 to 50 m in Madison Range and Gravelly Range, although is missing along east side of Gravelly Range (Hadley 1969b); thickness 50 to 75 m in northern Gallatin Range (McMannis and Chadwick, 1964).

Cambrian

- €rlp** **Red Lion Formation and Pilgrim Dolostone (Upper Cambrian, undivided) (€u of Kellogg and Williams, 2000)**—Mapped east of Gallatin River and north of Spanish Peaks fault (McMannis and Chadwick, 1964)
- €rpp** **Red Lion Formation and Pilgrim Dolostone (Upper Cambrian) and Park Shale (Middle Cambrian), undivided (€rp of Kellogg and Williams, 2000)**—Mapped south of Cedar Creek and east of Madison Valley (Tysdal, 1990; Kellogg, 1992).
- €pf** **Park through Flathead Formations, undivided (Middle Cambrian) (€mi of Kellogg and Williams, 2000)**—Mapped east of Gallatin River and north of Spanish Peaks fault (McMannis and Chadwick, 1964).
- €m** **Meagher Limestone (Middle Cambrian)**—Thin- to massive-bedded, light-gray to brownish-gray, finely crystalline limestone. Locally oolitic, especially in upper part. Upper 30 m is thin-bedded, gray limestone that contains conspicuous orange mottles; upper few meters contains fissile gray-green shale. Middle 50 m is medium- to massive-bedded limestone, locally mottled tan and containing small silicic limestone stringers. Lower 30 m is thin- to medium-bedded, tan-mottled, gray limestone that contains a few intercalated micaceous shale beds in lower part. Forms cliffs. Conformably overlies Wolsey Shale. About 100-150 m thick in Madison Range (Tysdal, 1990; Kellogg, 1992), 100-110 m thick in Gravelly Range (Hadley, 1980), but only about 50 m thick in northern Gallatin Range (McMannis and Chadwick, 1964).
- €wf** **Wolsey and Flathead Formations, undivided (Middle Cambrian)**
- Wolsey Shale:** Mostly thin-bedded, greenish-gray, olive-drab, gray, and grayish-brown micaceous sandstone, siltstone, and shale. Sandstone beds are wavy and bioturbated, contain green and gray- and green-mottled shale interbeds, and generally weather brown; animal trails are common; locally glauconitic. Near middle of unit is a 10- to 15-m-thick section of thin-bedded, dark-gray, brown-weathering argillaceous limestone interbedded with lesser amount of sandstone and shale. Upper 5 m is interbedded wavy-laminated, thin-bedded, gray limestone and gray, micaceous siltstone. Conformable with underlying Flathead Sandstone. Forms slopes and swales. Thickness 30-65 m.

Flathead Sandstone: Thin- to medium-bedded, medium- to coarse-grained, reddish-brown, tan, and purplish-tan, quartz-rich, feldspathic sandstone; locally weathers to rusty red. Two thin zones of fine-grained, micaceous, greenish-gray argillaceous sandstone near top of formation. Basal part of formation contains rounded pebbles of metamorphic rock. Unconformably overlies Archean crystalline rock. Thickness variable from 15 to 75 m.

Proterozoic and/or Archean

p \in db **Diabase dike (Proterozoic; ! di of Kellogg and Williams (2000))**—Black to dark-greenish-gray, fine- to medium-grained, equigranular, well indurated diabase in steep, northwest-striking dikes as wide as 30 m (most are considerably thinner). Contains about 30-50 Percent euhedral labradorite, 30-60 percent augite, 0-30 percent hornblende (inverted from augite), 0-10 percent potassium feldspar, 5-8 percent opaque minerals, 1-3 percent apatite, 0-5 percent biotite, 0-3 percent quartz, and from trace to 1 percent epidote. Emplaced during two periods: about 1,455 Ma and 750-780 Ma (Harlan and others, 1990).

Low grade metamorphic and intrusive rocks of the Ruby Creek Area, eastern Gravelly Range (Proterozoic or Archean)

p \in md **Metadiorite (! Amd of Kellogg and Williams, 2000)**—Mapped by Hadley (1969a, b) but not described in detail.

p \in ph **Phyllite (! Aph of Kellogg and Williams, 2000)**—Greenish-gray, greenish-red, gray-red, and dark-gray phyllite containing quartz, biotite, chlorite, muscovite, hematite, magnetite, and rare garnet. Contains thin layers of micaceous and actinolite-bearing quartzite and lenticular amphibolite bodies (not shown).

p \in em **Epidote-actinolite metasediment (! Ass of Kellogg and Williams, 2000)**—Green, metamorphosed, fine-grained clastic sedimentary rocks, locally showing good cross-bedding and graded beds. Contains quartz, plagioclase, actinolite, and epidote. Weathers to blocky outcrops.

p \in bs **Biotite-chlorite schist (! Abs of Kellogg and Williams, 2000)**—Dark-brownish-gray to black, well-foliated schist containing quartz, plagioclase, biotite, chlorite, magnetite, and muscovite; porphyroblasts of plagioclase abundant.

p \in if **Banded iron formation in Gravelly Range (! Aif of Kellogg and Williams, 2000)**—Alternating layers typically 2-10 cm thick, of reddish-brown to black magnetite-quartz and quartzite. Contains minor hydrobiotite and magnetite locally oxidized to hematite and goethite. Layers are commonly tightly folded.

p \in ms **Mylonitic schist (! Ams of Kellogg and Williams, 2000)**—Well-foliated chlorite-actinolite schist that contains abundant shear-bounded bodies of metagabbro, metadiorite, and amphibolite. Interpreted as sheared amphibolite (J.M. O'Neill, oral commun., 1995) and may be approximately equivalent to biotite-chlorite schist (unit p \in bs). Includes a thin staurolite-sillimanite mylonite that bounds southern contact of unit.

- p€gq** **Biotite gneiss, quartzite, and hornblende gneiss (! Agb of Kellogg and Williams, 2000)**—Approximately as mapped by Hadley (1969b); part of quartzfeldspar gneiss and pegmatite unit of Vargo (1990).
- p€gg** **Granite, migmatite, pegmatite, and granitic orthogneiss (! Agg of Kellogg and Williams, 2000)**—Approximately as mapped by Hadley (1969b), who did not describe unit in detail. Not known if unit is intrusive into adjacent units or forms older basement.

Archean

Metaigneous rocks

- Agp** **Granite porphyry of Hell Roaring Creek**—Pink, coarse-grained, massive to slightly foliated biotite monzogranite porphyry. Contains about 20 percent conspicuous euhedral to subhedral potassium-feldspar phenocrysts as long as 1.5 cm; matrix contains about 30 percent quartz, 35 percent plagioclase, 15 percent potassium feldspar and about 20 percent biotite. Locally sheared, forming well-developed pinkish-gray augen gneiss, as on Indian Ridge in the Spanish Peaks.
- Agog** **Granitic orthogneiss (Agg of Kellogg and Williams, 2000)**—Light-gray to light-pinkish-gray, generally tan-weathering, medium-grained, hypidiomorphic, weakly to moderately foliated orthogneiss generally ranging in composition from tonalite to monzogranite. Mafic mineral almost exclusively biotite (trace to 15 percent): may contain as much as 5 percent almandine and, rarely, 5 percent hornblende, 2 percent augite, and 2 percent muscovite; also contains traces of zircon, epidote, allanite, and opaque minerals. One unusual occurrence is a pluton near Summit Lake in the Spanish Peaks; the pluton is a discordant body composed of massive to weakly foliated, very light gray biotite tonalite.
- Amb** **Metabasite**—Black (commonly speckled with white feldspar and pink garnet), fine-grained, equigranular, granoblastic, weakly foliated to massive hornblende-augite-almandine metabasite. Composition variable; contains 15-45 percent plagioclase (mostly andesine), 10-60 percent yellowish-green to brown hornblende, 2-20 percent augite, 0-20 percent almandine, 0-5 percent reddish-brown biotite, 1-3 percent opaque minerals, and trace of apatite. In some places relict porphyritic texture is preserved as white clusters of fine-grained plagioclase crystals as long as 1 cm. Mostly occurs as sills as wide as about 50 m concordant to foliation; in some places sills show pinch-and-swell structure and boudinage. Commonly enveloped in medium-grained amphibolite margins as wide as 10 m, indicating post-emplacement metasomatism at amphibolite grade. Equivalent to orthoamphibolite of Vitaliano and Cordua (1979) in Tobacco Root Mountains.
- Ahgs** **Hornblende-biotite granodiorite orthogneiss of Summit Lake (Ags of Kellogg and Williams, 2000)**—Light- to medium-gray, medium-grained, hypidiomorphic, poorly to well foliated hornblende-biotite granodiorite orthogneiss. Contains 30-40 percent plagioclase, 10-20 percent potassium-feldspar, 15-25 percent quartz, 10-20 percent hornblende, and about 10 percent biotite. Most rock is highly strained, forming well developed foliation; in such places, mafic minerals are concentrated in ribbon layers 2-5 mm thick. Contains numerous, thin (3-

10 mm) feldspar-quartz migmatitic layers. Cut by at least three periods of sill and dike intrusion; sills and dikes locally so closely spaced that they form an agmatite. Cut by granitic orthogneiss of Summit Lake plutons.

Amum **Meta-ultramafic rocks (Aum of Kellogg and Williams, 2000)**—Black to dark-greenish-gray, fine- to medium-grained, well-foliated to massive, variably serpentinized ultramafic rocks of wide-ranging composition; includes olivine websterite, iherzolite, and olivine clinopyroxenite. Accessory minerals include olive-green spinel, magnetite, and apatite. Commonly contains secondary amphibolite (anthophyllite or actinolite) serpentine, talc, dolomite, magnesite, and (or) mica. Occurs in lenses, pods, and small irregularly shaped masses, rarely more than 10 m in diameter. Probably tectonically incorporated into country rock.

Gneissic rocks of unknown origin (Archean)

Aqfg **Quartzofeldspathic gneiss (Aqf of Kellogg and Williams, 2000)**—Mapped where rocks are heterogeneous, generally layered, light- to medium-gray microcline-plagioclase-quartz-biotite gneiss. Commonly migmatitic and blastomylonitic. Some areas not mapped in detail, such as south of Sphinx Mountain in Madison Range and east of Gallatin River in Gallatin Range, which may contain many of the other gneissic rock units.

Agph **Garnetiferous gneiss of the Tobacco Root Mountains (Agn of Kellogg and Williams, 2000)**—Includes biotite-garnet schist, sillimanite-garnet schist, garnetiferous quartzite, quartzite, highly garnetiferous quartzofeldspathic gneiss, corundum gneiss, gedrite schist, cummingtonite schist, and garnetiferous amphibolite (Vitaliano and Cordua, 1979). May represent refractory component (“restite”) remaining after removed of partial melts.

Aamh **Amphibolite and hornblende-plagioclase gneiss (Aam of Kellogg and Williams, 2000)**—Most is gray to black, medium-grained, hypidiomorphic equigranular, moderately foliated to well-foliated hornblende-plagioclase gneiss and amphibolite; contains as much as 5 percent quartz and traces of zircon, opaque minerals, and apatite; locally garnetiferous. Plagioclase is typically An₃₀ and weathers white. Commonly contains white, migmatitic leucosomes of anorthosite as thick as 10 cm. Similar unit in tobacco Root Mountains interpreted to be either metamorphosed, clay-rich dolomite or mafic-extrusive rock (Vitaliano and Cordua, 1979). Amphibolite envelopes around some metabasite intrusive bodies indicate at least some amphibolite was derived from intrusive rocks. Unit may include minor amounts of other Archean units.

Abs **Biotite schist**—Black, dark-gray, and gray, fine- to medium-grained biotite-plagioclase-quartz, ±hornblende, ±microcline schist. Interpreted to be sheared mafic rock that equilibrated at lower amphibolite facies; not studied in detail. Crops out in Gallatin Canyon-Hell Roaring Lake (Spanish Peaks) area.

Agd **Gedrite and cummingtonite gneiss**—Cummingtonite gneiss is light gray, medium grained, strongly lineated, and vitreous. Contains as much as 90 percent cummingtonite, and variable amounts of quartz, almandine, muscovite, rutile, and opaque minerals. Grades into metaquartzite. Gedrite gneiss is brown to grayish-brown, moderately well foliated gneiss that contains as much as 70 percent clove-brown gedrite, in addition to quartz, plagioclase ±

sillimanite, ±biotite, ±cordierite, and traces of magnetite and rutile. Both rock types occur as small lenses and concordant layers in other Archean rocks.

Abhg **Biotite-hornblende gneiss of Beartrap Canyon (Abh of Kellogg and Williams, 2000)**—White, light-gray, dark-gray, and black, medium-grained, well-foliated and well-layered gneiss. Leucosomes contain plagioclase, quartz, biotite, ±potassium feldspar, ±garnet, and a trace of opaque minerals. Melasomes contain biotite, hornblende, plagioclase, quartz, ±garnet, and trace of opaque minerals. Contains rare quartzite layers and sillimanite-bearing gneiss. Commonly migmatitic. Layers are typically 1-20 cm thick. Injected by numerous sills of metabasite. Crops out extensively in Bear Trap Canyon of Madison River. Similar rocks, also mapped as unit Abh [Abhg], occur at several other less extensive localities.

Aamh* **Amphibolite and hornblende gneiss (from central border with Bozeman quadrangle)***

High-grade tectonites (Archean)

Amyc **Mylonite of Crooked Creek shear zone (At of Kellogg and Williams, 2000)**—Light- to medium-gray, mottled mylonite that has a well-developed L-S tectonite fabric and a fine-grained recrystallized (granoblastic) texture. Composition variable, ranging from that of quartzofeldspathic gneiss to amphibolite. The mylonite equilibrated at near-peak (lower granulite) metamorphic conditions and forms anastomosing, concordant zones in northern Madison Range (Kellogg, 1993a; Kellogg and Mogk, 1991). Margins of unit are concordant with, and grade, into relatively unmylonitized gneiss.

Metasedimentary rocks (Archean)

Aags **Aluminous gneiss and schist (Aag of Kellogg and Williams, 2000)**—Gray to dark-brownish-gray, medium-grained, inequigranular, generally well foliated, commonly micaceous gneiss and schist containing aluminosilicate (mostly sillimanite and rarer kyanite). Unit contains 5-90 percent anhedral quartz having undulatory extinction, 0-30 percent microcline, 0-35 percent plagioclase, 0-30 percent almandine, 0-20 percent muscovite, tract to 15 percent sillimanite or kyanite, 0-10 percent reddish-brown biotite, 0-2 percent opaque minerals, including graphite, and trace of zircon. Commonly rich in quartz and locally grades into quartzite. Several kyanite prospects are on east side of Gravelly Range, in sec. 6, T. 8 S., R. 1 W. (Nordstrom, 1947).

Abmg **Biotite-muscovite gneiss (Abm of Kellogg and Williams, 2000)**—Light- to medium-gray, medium-grained, poorly foliated to well-foliated quartz-feldspar gneiss that contains abundant thin schistose layers containing both biotite and muscovite. Aluminosilicate-bearing lenses are common, and one prominent 20-to 50-m-thick schistose horizon contains as much as 50 percent coarse-grained biotite, about 20 percent quartz, 0-10 percent gedrite, 5 percent plagioclase, 5 percent sillimanite, 3 percent kyanite (as large blue blades), 3 percent garnet, and 3 percent muscovite. Mapped on north side of Hell Roaring Creek Valley in Spanish Peaks.

Aq **Quartzite**—White, medium- to coarse-grained inequigranular, moderately foliated to massive quartzite; locally bright green where trace amounts of chromium-bearing mica (fuchsite) are present. In most places unit is composed entirely of anhedral quartz grains having undulatory

extinction, but locally contains as much as 30 percent microcline, 20 percent muscovite, 15 percent sillimanite, 10 percent cummingtonite, 8 percent almandine, 2 percent actinolite, and trace of zircon and opaque minerals. Commonly forms prominent ridges, especially in northwestern part of Madison Range, where core of a plunging, nearly isoclinal antiform is composed almost entirely of quartzite. Unit is interlayered in most places with mafic amphibolite.

Am **Marble (Ama of Kellogg and Williams, 2000)**—White, coarse-grained massive to moderately well foliated dolomitic marble that contains as much as 3 percent quartz grains. Weathers orange-brown. Locally hydrothermally altered to commercial deposits of talc on east side of Gravelly Range. A few thin layers of calcitic marble, containing as much as 30 percent serpentine, mapped in northern part of Madison Range. Extensive, commercially exploited talc deposits occur in hydrothermally altered zones in large marble body on eastern side of Gravelly Range.

Aif **Banded iron formation**—Magnetite-bearing schist and gneiss, gruehnite-magnetite schist, and garnet-quartz rock. Mapped in Tobacco Root Mountains (Vitaliano and Cordua, 1979).

GEOLOGIC MAP SYMBOLS

Ennis 30' x 60' Quadrangle



Contact: Dashed where approximately located. Dotted where concealed. Queried where uncertain



Normal fault: Dashed where approximately located, dotted where concealed. Queried where uncertain. Bar and ball on downthrown side where known.



Thrust fault: Dashed where approximately located, dotted where concealed, teeth on upthrown block



Reverse fault: Dashed where approximately located, dotted where concealed, open teeth on upthrown block



Low angle normal fault: Dashed where approximately located, dotted where concealed.



Anticline: Showing trace of axial plane and direction of plunge, dotted where concealed



Overturned anticline: Showing trace of axial plane and direction of dip of bedding; dashed where approximately located, dotted where concealed



Syncline: Showing trace of axial plane and direction of plunge, dotted where concealed



Overturned syncline: Showing trace of axial plane and direction of dip of bedding; dashed where approximately located, dotted where concealed



Strike and dip of inclined beds



Strike and dip of vertical beds



Strike and dip of overturned beds



Horizontal beds



Strike and dip of inclined foliation



Strike and dip of vertical foliation

References

Primary reference for this quadrangle map:

Kellogg, K.S. and Williams, V.S., 2000, Geologic map of the Ennis 30'x 60' quadrangle Madison and Gallatin Counties, Montana, and Park County, Wyoming: U.S. Geological Survey Geologic Investigation Series I-2690, 1:100,000 scale.

References from Kellogg and Williams (2000)

Bearzi, J.P., 1987, Soil development, morphometry, and scarp morphology of fluvial terraces at Jack Creek, southwestern Montana: Bozeman, Montana State University, M.S. thesis, 131 p.

Chadwick, R.A., 1969, The northern Gallatin Range, Montana—northwestern part of the Absaroka-Gallatin volcanic field: University of Wyoming Contributions to Geology, v. 8, no. 2, pt. 2, p. 150–156

Christiansen, R.L., and Blank, H.R., Jr., 1972, Volcanic Stratigraphy of the Quaternary rhyolite plateau in Yellowstone National Park: U.S. Geological Survey Professional Paper 729–B, p. B1–B18.

DeCelles, D.G., and 15 others, 1987, Laramide thrust-generated alluvial-fan sedimentation, Sphinx conglomerate, southwestern Montana: American Association of Petroleum Geologists Bulletin, v. 71, p. 135–155.

Hadley, J.B., 1969a, Geologic map of the Cameron quadrangle, Madison County, Montana: U.S. Geological Survey Geologic Quadrangle Map GQ–813, scale 1:62,500.

Hadley, J.B., 1969b, Geologic map of the Varney quadrangle, Madison County, Montana: U.S. Geological Survey Geologic Quadrangle Map GQ–814, scale 1:62,500.

Hadley, J.B., 1980, Geology of the Varney and Cameron quadrangles: U.S. Geological Survey Bulletin 1459, 108 p.

Hall, W.B., 1961, Geology of part of the upper Gallatin Valley of southwestern Montana: Laramie, University of Wyoming, Ph.D. thesis, 239 p.

Hanneman, D.L., and Wideman, C.J., 1991, Sequence Stratigraphy of Cenozoic continental rocks, southwestern Montana: Geological Society of American Bulletin, v. 103, p. 1335–1345.

Hanson, A.M., 1952, Cambrian stratigraphy in southwestern Montana: Montana Bureau of Mines and Geology Memoir 33, 46 p.

Harlan, S.S., Snee, L.W., and Geissman, J.W., 1990, Paleomagnetic and $^{40}\text{Ar}/^{39}\text{Ar}$ results from mafic dikes and basement rocks, southwest Montana: EOS, Transactions of the American Geophysical Union, V. 71, p. 1297.

- Hiza, M.M., 1994, Processes of alluvial sedimentation in Eocene Hyalite Peak volcanics, Absaroka-Gallatin volcanic province, southwest Montana: Bozeman, Montana State University, M.S. thesis, 163 p.
- Kellogg, K.S., 1992, Geologic map of the Fan Mountain quadrangle, Madison County, Montana: U.S. Geological Survey Geologic Quadrangle Map GQ-1706, scale 1: 24,000.
- Kellogg, K.S., 1993a, Geologic map of the Cherry Lake quadrangle, Madison County, Montana, U.S. Geological Survey Geologic Quadrangle Map GQ-1725, scale 1: 24, 000.
- Kellogg, K.S., 1993b, Geologic map of the Ennis Lake quadrangle, Madison County, Montana: U.S. Geological Survey Geologic Quadrangle Map GQ-1729, scale 1:24, 000.
- Kellogg, K.S., 1994, Geologic map of the Norris quadrangle, Madison County, Montana: U.S. Geological Survey Geologic Quadrangle Map GQ-1738, scale 1: 24,000.
- Kellogg, K.S., 1995, Geologic map of the Bear Trap Creek quadrangle, Madison County, Montana: U.S. Geological Survey Geologic Quadrangle Map GQ-1757, scale 1: 24,000.
- Kellogg, K.S., and Mogk, D.W., 1991, The Crooked Creek mylonite—A high-temperature zone of ductile deformation in the Archean of the northern Madison Range, southwestern Montana [abs]: Geological Society of America Abstracts with Programs, v. 23, no. 5, p. A 59.
- Kellogg, K.S., Schmidt, C.J., and Young, S.W., 1995, Basement and cover-rock deformation during Laramide contraction in the northern Madison Range (Montana) and its influence on Cenozoic basin formation: American Association of Petroleum Geologists Bulletin, v 79, p. 1117–1137.
- Lauer, T.C., 1967, Stratigraphy and structure of the Snowflake Ridge area, Gallatin County, Montana: Corvallis, Oregon State University, M.S. thesis, 182 p.
- Marvin, R.F., Wier, K.L., Mehnert, H.H., and Merritt, V.M., 1974, K-Ar ages of selected Tertiary igneous rocks in southwestern Montana: Isochron/West, no. 10, p. 17–20.
- McMannis W.J., and Chadwick R.A., 1964, Geology of the Garnet Mountain quadrangle, Gallatin County, Montana: Montana Bureau of Mines and Geology Bulletin 43, 47 p.
- Nordstrom, C. L., 1947, Geology of a kyanite deposit near Ennis, Montana: Butte, Montana School of Mines, Senior thesis, 30 p.
- Rasmussen D.L., and Fields, R.W., 1983, Structural and depositional history, Jefferson and Madison basins, southwestern Montana [abs.]: American Association of Petroleum Geologists Bulletin, v. 67, p. 1352
- Ritter, J.B., and 16 others, 1990, The Late Quaternary geology of Cedar Creek alluvial fan, Madison River Valley, southwestern Montana, *in* Hall, R.D., ed., Quaternary geology of the western Madison Range, Madison Valley, Tobacco Root Range, and Jefferson Valley; Rocky Mountain Friends of the Pleistocene Fieldtrip Guidebook: Indianapolis, Indiana University, p. 120-138.
- Salt, K.L., 1987, Archean geology of the Spanish Peaks area, southwestern Montana: Bozeman, Montana State University, M.S. thesis, 81 p.

- Schneider, N.P., and Ritter, D.F., 1987, Late Pleistocene response of the Madison River to regional base level change for the Madison Valley, southwest Montana [abs.]: Geological Society of America Abstracts with Programs, v. 19, no. 5, p. 332.
- Simons, F.S., Van Loenen, R.E., and Moore, S.L., 1985, Geologic map of the Gallatin Divide roadless area, Gallatin and Park Counties, Montana: U.S. Geological Survey Miscellaneous Field Studies Map MF-1569-B, scale 1:62,500.
- Smedes, H.W., and Prostka, H.J., 1972, Stratigraphic framework of the Absaroka Volcanic Supergroup in the Yellowstone National Park region: U.S. Geological Survey Professional Paper 729-C, p. C1-C33.
- Streckeisen, Albert, 1976, To each plutonic rock its proper name: Earth-Science Reviews, v. 12, p. 1-33.
- Swanson, R.W., 1950, Geology of part of the Virginia City and Eldridge quadrangles, Montana: U.S. Geological Survey Open-File Report 51-4, 12 p.
- Todd, S.G., 1969, Bedrock geology of the southern part of Tom Minor Basin, Park and Gallatin Counties, Montana: Bozeman, Montana State University, MS. thesis, 62 p.
- Turner, F.J., and Verhoogen, John, 1960, Igneous and metamorphic petrology: McGraw-Hill Book Company, Inc., New York, 694 p.
- Tysdal, R.G., 1990, Geologic map of the Sphinx Mountain quadrangle and adjacent parts of the Cameron, Cliff Lake, and Hebgen Dam quadrangles, Montana: U.S. Geological Survey Miscellaneous Investigations Series Map I-1815, scale 1:62,500.
- Tysdal, R.G., Marvin, R.F., DeWitt, E.H., 1986, Late Cretaceous stratigraphy, deformation, and intrusion in the Madison Range of southwestern Montana: Geological Society of America Bulletin, v. 97, p. 859-868.
- Tysdal, R.G., Simons, F.S., 1985, Geologic map of the Madison Roadless Area. Gallatin and Madison counties, Montana: U.S. Geological Survey Miscellaneous Field Studies Map MF-1605-B, scale 1:96,000.
- U.S. Geological Survey, 1972, Geologic map of Yellowstone National Park: U.S. Geological Survey Miscellaneous Investigations Map I-711, scale 1:125,000.
- Vargo, A.G., 1990, Structure and petrography of the pre-Beltian rocks of the north-central Gravelly Range, Montana: Fort Collins, Colorado State University, M.S. thesis, 157 p.
- Vitaliano, C.J., and Cordua, W.S., 1979, Geologic map of southern Tobacco Root Mountains, Madison County, Montana: Geological Society of America Map and Chart Series MC-31, scale 1:62,500.
- Wardlaw, B.R., and Pecora, W.C., 1985, New Mississippian-Pennsylvanian stratigraphic units in southwest Montana and adjacent Idaho: U.S. Geological Survey Bulletin 1656, p. B1-B9.
- Wier, K.L., 1982, Map showing geology and outcrops of the Virginia City and Alder quadrangles, Madison County, Montana: U.S. Geological Survey Miscellaneous Field Studies Map MF-1490, scale 1:12,000.

Additional references


Berg, R.B., Lonn, J.D., and Locke, W.W., 1999m Geologic map of the Gardiner 30' x 60' quadrangle, south-central Montana: Montana Bureau of Mines and Geology Open File Report 387, scale 1:100,000.

Berg, R.B., Lopez, D.A., and Lonn, J.D., 2000, Geologic map of the Livingston 30' x 60' quadrangle, south-central Montana: Montana Bureau of Mines and Geology Open File Report 406, scale 1:100,000.

O'Neill, J.M., and Christiansen, R.L., 2001, Preliminary geologic map of the Hebgen Lake 30' x 60' quadrangle, Montana, Wyoming, and Idaho: Montana Bureau of Mines and Geology Open File Report 464, scale 1:100,000. Later published as O'Neill, J.M., and Christiansen, R.L., 2004, Geologic map of the Hebgen Lake quadrangle, Beaverhead, Madison, and Gallatin Counties, Montana, Park, and Teton Counties, Wyoming, and Clark and Freemont Counties, Idaho: U. S. Geological Survey Scientific Investigations Map 2816, scale 1:100,000.

Vuke, S.M., Lonn, J.D., Berg, R.B., and Kellogg K.S., 2002, Preliminary geologic map of the Bozeman 30' x 60' quadrangle, southwestern Montana: Montana Bureau of Mines and Geology Open File Report 469, scale 1:100,000.

Ground Water Data



Groundwater Information Center
Montana Bureau of Mines and Geology
Montana Tech of The University of Montana
 1300 West Park Street - Main Hall 314
 Butte Montana 59701-8997

You are currently signed in. | 4/29/2009
[Sign Out](#)

[Home](#) | [Well Data](#) | [Reports](#) | [Data Coop](#) | [DrillerWeb](#) | [DNRC](#) | [Help!](#)

MbmgGwic Navigation: | [Main Menu](#) | [SWL Menu](#) | [GWCP Map Products](#) | [Ground-Water Projects](#) |

GWIC Data > Well Construction Data > Township: 04S Range: 02W Sec: 26, 35

The following data were returned from the GWIC databases for the area you requested. For a more detailed description of the data view the [GWIC Metadata report](#). If you notice data entry errors or have questions please let us know by clicking [here](#) to leave us a message. If you wish to view a one page report for a particular site, click the hyperlinked **Gwic Id** for that well. Scroll to the right of your screen to view all the data. All data displayed on the screen may not show up when printed.

Retrieval Statistics*

Field	Max	Min	Avg
Total Depth (ft)	400.00	40.00	197.76
Static Water Level (ft)	230.00	18.00	72.94
Yield (gpm)	300.00	10.00	39.47

* These statistics do not take any geographic, topographic, or geologic factors into consideration. Negative swl values are reported for water levels that are above land surface.

Did you know about...

Other GWIC data

GWIC has 1 water quality sample(s) for this area.
GWIC has 1 field visit(s) for this request area.
GWIC has 13 water level(s) for this request area.

Thanks, Just take me back to the menu.

Other MBMG data

MBMG has 439 publications available for MADISON county.
MBMG has 12 abandoned mine record(s) for this request area.

Gwic Id	PDF	DNRC WR	Site Name	Twn	Rng	Sec	Q Sec	Ver?	Type	Td	Swl	Pwl	Rwl	Yield	Test	Date	Use
142705			MURPHY JAN	04S	02W	26		No	WELL	325.00	96.00	320.00		12.00	AIR	12/6/1993	DOMESTIC
165088		C098022-00	BORSUK JEFF	04S	02W	26	BCA	No	WELL	285.00	113.00		113.00	20.00	AIR	9/5/1995	DOMESTIC
142706		C097920-00	WOODS JAMES B.	04S	02W	26	BDC	No	WELL	200.00	116.00	195.00		15.00	AIR	10/4/1993	DOMESTIC
107775		C30042942	FENDRICK CARL	04S	02W	26	CA	No	WELL	260.00	177.00	192.00		20.00	OTHER	10/1/1982	UNKNOWN
107776			HALL EMERSON	04S	02W	26	CBB	No	WELL	400.00				10.00	OTHER	8/23/1982	DOMESTIC
130120		C082669-00	MARTIN DR JAMES L	04S	02W	26	CBB	No	WELL	287.00	230.00	280.00		15.00	AIR	3/24/1992	DOMESTIC
126652			SCHAUWECKER THOMAS	04S	02W	26	CC	No	WELL	103.00	32.00	98.00		15.00	AIR	11/7/1991	DOMESTIC
228605		C30024941	STEPHENS, RICHARD AND BARBARA	04S	02W	26	CCDD	No	WELL	280.00	45.00		45.00	35.00	AIR	8/15/2006	DOMESTIC
142632			GOUGH STEPHEN OR GALE	04S	02W	35		No	WELL	145.00	82.00	140.00	82.00	15.00	AIR	5/17/1994	DOMESTIC
204531			HOE TIM AND RENE	04S	02W	35		No	WELL	100.00	48.00		48.00	75.00	AIR	7/9/2003	DOMESTIC
210381			RAMSHORN RANCH LLC	04S	02W	35		No	WELL	160.00	40.00		40.00	300.00	AIR	3/11/2004	STOCKWATER
216049			RAMSHORN RANCH LLC	04S	02W	35		No	WELL	40.00	23.00		23.00	10.00	AIR	10/22/2004	STOCKWATER
230286		C30027908	GAAR GILL	04S	02W	35	ABCB	No	WELL	145.00	30.00			35.00	AIR	9/20/2006	DOMESTIC
167346			MILLER ROBERT	04S	02W	35	ADD	No	WELL	85.00	28.00		28.00	25.00	AIR	2/9/1998	DOMESTIC
140954		C088045-00	LEADBEATER NANCY	04S	02W	35	BA	No	WELL	200.00	31.00		32.00	12.00	AIR	9/14/1993	DOMESTIC
221978			PATRICK JAMES AND JANE	04S	02W	35	BA	No	WELL	100.00	18.00		18.00	45.00	AIR	7/8/2005	DOMESTIC
120760		C077940-00	GOODMAN JEFF AND CINDY SHARP	04S	02W	35	BACB	Yes	WELL	247.00	58.00			12.00	AIR	6/27/1990	DOMESTIC

End of Report.
 17 record(s) listed.

Items of Note:

¹This report is restricted to site types of WELL, BOREHOLE, SPRING, COAL BED METHANE WELL, PETWELL, PIEZOMETER.






²A single well record (a distinct GWIC Id) may be represented by more than one line in this report if more than one performance test was conducted on the well at

the time of drilling.

Explanation of Columns:

GWIC Id = Key field for the GWIC database. Links to one page reports.

PDF = Are scanned documents available through the Document Manager?

-  = Yes, click on the icon to download the PDF file.
-  = No, well was submitted electronically. No paper record exists.
-  = No, record does have a known well log but it is not scanned yet.
-  = No, record may or may not have a document to scan. Metadata is unclear.
-  = No, record was created from a source other than a well log. No paper record exists.

DNRC WR = Water right number assigned to this site by Department of Natural Resources and Conservation.

Site Name = Current owner name assigned to GWIC record.

Location = Location of site in Montana township, range, section, and quarter-section coordinates.

Ver? = Has this location been verified by field staff?

Type = Type of site assigned to GWIC record.

Td = Total depth of well in feet below ground.

Swl = Static water level in feet above/below ground - Negative values are reported for water levels that are above land surface.

Pwl = Pumping water level in feet below ground.

Rwl = Recovery water level in feet below ground.

Yield = Yield in gallons per minute.

Test = Type of performance test reported.

Date = Completion date of well/borehole.

Use = Reported use of water.

Disclaimer:

The preceding materials represent the contents of the GWIC databases at the Montana Bureau of Mines and Geology at the time and date of the retrieval. The information is considered unpublished and is subject to correction and review on a daily basis. The Bureau warrants the accurate transmission of the data to the original end user at the time and date of the retrieval. Retransmission of the data to other users is discouraged and the Bureau claims no responsibility if the material is retransmitted. There may be wells in the request area that are not recorded at the Information Center.

Ground-Water Information Center Online © 1998 - 2009

[Staff](#) | [Privacy Statement](#)